




Contents lists available at ScienceDirect

Journal of Molecular Liquids

journal homepage: www.elsevier.com/locate/molliq



Corrigendum

Corrigendum for development of Abraham model correlations for solute transfer into 2-methoxyethanol from water and from the gas phase at 298 K [J. Mol. Liq. (2015), 209, 738–744]



Erin Hart^a, Damini Grover^a, Heidi Zettl^a, Victoria Koshevarova^a, Shoshana Zhang^a, Colleen Dai^a, William E. Acree Jr.^{a,*}, Igor A. Sedov^b, Mikhail A. Stolov^b, Michael H. Abraham^c

^a Department of Chemistry, University of North Texas, 1155 Union Circle Drive #305070, Denton, TX 76203, (USA)

^b Department of Chemistry, Kazan Federal University, Kremlevskaya 18, Kazan 420,008, Russia

^c Department of Chemistry, University College London, 20 Gordon Street, London WC1H 0AJ, UK

ARTICLE INFO

Article history:

Received 19 February 2017

Accepted 21 February 2017

Available online 28 February 2017

© 2017 Elsevier B.V. All rights reserved.

The mole fraction solubility of trans-stilbene in 2-methoxyethanol was inadvertently given in Table 1 as $X_{S,organic}^{exp} = 0.1442$. The correct value is $X_{S,organic}^{exp} = 0.01442$.

DOI of original article: <http://dx.doi.org/10.1016/j.molliq.2015.05.037>.

* Corresponding author.

E-mail address: acree@unt.edu (W.E. Acree).